

WAVELET PERSPECTIVE OF A DISORIENTED CHIRAL CONDENSATE ^a

Zheng Huang[†], Ina Sarcevic[†], Robert Thews[†] and Xin-Nian Wang[‡]

[†]*Department of Physics, University of Arizona, Tucson, AZ 85721*

[‡]*Nuclear Science Division, LBNL, Berkeley, CA 94720*

The possible chiral phase transition in high energy heavy-ion collisions may lead to the formation of a disoriented chiral condensate (DCC). However, the existence of many uncorrelated small domains in the rather large interaction volume and the huge combinatorial background make the experimental search for the DCC signal a rather difficult task. We present a novel method for studying the formation of a DCC in high energy heavy-ion collisions utilizing a discrete wavelet transformation. We find the wavelet power spectrum for a DCC to exhibit a strong dependence on the scale while an equilibrium system and the standard dynamical models such as HIJING have a flat spectrum.

In heavy-ion collisions the rapid expansion of the collision debris in the longitudinal direction may lead to supercooling of the interaction region, and as a result, domains of the “unconventionally” oriented vacuum configurations allowed by the chiral symmetry may be formed.¹ Detection of this interesting phenomenon, the so-called Disoriented Chiral Condensate (DCC)¹, would provide valuable information on the vacuum structure of the strong interaction and the nature of the chiral phase transition. Preliminary theoretical investigations on non-equilibrium dynamics using the classical linear σ -model have found some evidence for the growth of long wavelength pion modes, which may indeed lead to domain structure or cluster formation.^{2,3} If there are many uncorrelated small domains, the integrated probability distribution of the neutral pion fraction f emitted from a disoriented region, predicted to be $P(f) = 1/2\sqrt{f}$,¹ would become Gaussian. This makes the experimental search for the DCC signal a rather difficult task. In order to disentangle the DCC domain structure in high energy heavy-ion collisions, we propose a new method which emphasizes not only the behavior of the probability distribution in the full phase space region but also its fluctuation in rapidity η or azimuthal angle ϕ . In other words, one needs to study the “local” properties of the distribution in phase space if the DCC clusters are “localized” objects in phase space.

A novel method is a multi-resolution analysis performed by a discrete wavelet transformation (DWT)⁴ which has been found effective in systematically detecting structures of various scales in turbulence, and multiparticle

^aTalk presented by Ina Sarcevic at DPF '96 and by Zheng Huang at PANIC '96

productions^{5,6} For any one-dimensional sample of a point distribution $f(x)$, such as the rapidity distribution of the neutral pion fraction in the interval $[0, 1]$ with resolution Δx , can be represented as a histogram of 2^J bins where $J = \text{mod}(|\ln \Delta x|/\ln 2) + 1$

$$f(x) \equiv f^{(J)}(x) = \sum_{k=0}^{2^J-1} f_{Jk} \phi_{Jk}(x), \quad (1)$$

where $\phi_{Jk}(x)$ are the piecewise constant functions called the mother functions. The histogram function at the next finer resolution is the average of adjacent bins at the finest scale. Evidently some detail is lost compared to (1). The difference between (1) and $f^{(J-1)}$ can be fully expressed in terms of the difference functions $\psi_{J-1,k}(x)$ which are stairwell functions called father functions. The difference amplitudes or one defines the DWT coefficients

$$\tilde{f}_{jk} = 2^j \int f(x) \psi_{jk}(x) dx, \quad (2)$$

where the basis functions $\psi_{jk}(x)$ have a unique property: they are related to each other by scaling and translation: $\psi_{jk}(x) = \psi(2^j x - k)$. The index j denotes the resolution scale and k the position at each scale. The idea of multiresolution analysis is to find representations of the sample function $f(x)$ at various scales. Unlike the Fourier transformation which requires the information in whole physical space, the wavelet SSD requires only the local information in space. Roughly speaking, the multiresolution analysis is a journey down (or up) a hierarchy of scales and to view the “world” in different “eyes” with magnifying or reducing glasses.

The central idea of using the wavelet analysis is to attempt an unbiased separation of the DCC physics from the various “backgrounds” which cannot be otherwise easily eliminated. The usual correlation method is not so effective here because the different scale correlations are mixed together in the same phase space. The discrete wavelet transformation (DWT), on the other hand, has an extra dimension – the scale which can be used to separate these correlations. The main source of the “backgrounds” is the fluctuations at small correlation scale due to pion radiation from the uncorrelated regions in space-time, which can coexist with a DCC domain. The background can also come from the partial thermalization of the system leading to thermal correlations. We have calculated the wavelet power spectrum for rapidity distribution of the neutral pion fraction simulated using the classical linear σ model.³ One defines




Figure 1: The power spectra of the wavelet coefficients for different dynamical scenarios.

the power spectrum with respect to the wavelet basis at each j as

$$P_j = \frac{1}{2^j} \sum_{k=0}^{2^j-1} |\tilde{f}_{jk}|^2. \quad (3)$$

P_j is then the power of fluctuations on length scale $L/2^j$. It is also the mean square of the probability distributions of the wavelet coefficients defined in (2). The result is shown in Fig.1. For the random samples and the equilibrium evolution of the classical field where there is no DCC production mechanism, the wavelet power is flat indicating there is no sensitivity on the change of the scale. For the DCC samples, the power starts to build up as j goes down below $j_d \sim 4$ and rises rather quickly, departing distinctively from the equilibrium and the random cases. This suggests the existence of the DCC clustering with a typical size of $\Delta\eta_d = 2\eta_{\max}/2^{j_d} \simeq 0.8$ units in rapidity. Also plotted is the power spectrum from HIJING Monte Carlo data which also features a flat distribution.

1. J. -P. Blaizot and A. Krzywcki, Phys. Rev. **D46**, 246 (1992); J. D. Bjorken *et al.*, SLAC preprint SLAC-PUB-6109.
2. K. Rajagopal and F. Wilczek, Nucl. Phys. **B404**, 577 (1993).
3. Z. Huang and X.-N. Wang, Phys. Rev. **D49**, R4339 (1994).
4. Z. Huang, I. Sarcevic, R. Thews and X.-N. Wang, Phys. Rev. **D54**, 750 (1996).
5. M. Farge, Ann. Rev. Fluid. Mech. **24**, 395 (1992).
6. M. Greiner, P. Lipa and P. Carruthers, Phys. Rev. **E51**, 1948 (1995)